



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

THE PERMANENT GRAVITATIONAL FIELD IN THE EINSTEIN THEORY.

BY LUTHER PFAHLER EISENHART.

1. In accordance with the theory of Einstein a permanent gravitational field is defined by a quadratic differential form

$$(1) \quad ds^2 = \sum_{i,k}^{1,\dots,4} g_{ik} dx_i dx_k \quad (g_{ik} = g_{ki}),$$

where the g 's, called the potentials of the field, are determined by the condition of satisfying ten partial differential equations of the second order, $G_{ik} = 0$. When the four coördinates x_i are functions of a single parameter, the locus of the point with these coördinates is a curve in four-space. If these functions are of such a character that the integral

$$(2) \quad \int \sqrt{\sum g_{ik} dx_i dx_k}$$

is stationary along the curve, the curve is called a "world-line," or a geodesic, in the four-space.

Einstein* considered the case when x_1, x_2, x_3 , are rectangular coördinates and x_4 represents the time, and assumed that the field was produced by a mass at the origin which did not vary with the time. In order to obtain the equations of the world-lines in the form which enabled him to establish his well-known expression for the precession of the perihelion of Mercury, Einstein made also the following assumptions:

- A. The quantities g are independent of t .
- B. The equations $g_{i4} = 0$ for $i = 1, 2, 3$.
- C. The solution is spacially symmetric with respect to the origin of coördinates in the sense that the solution is unaltered by an orthogonal transformation of x_1, x_2, x_3 .
- D. At infinity the quantities $g_{ik} = 0$ for $i \neq k$, and

$$g_{44} = -g_{11} = -g_{22} = -g_{33} = 1.$$

Schwarzschild† using the first three of these assumptions and certain others integrated the equations $G_{ik} = 0$, and obtained (1) in the form

* Berlin Sitzungsberichte, 1915, p. 831.

† Berlin Sitzungsberichte, 1916, p. 189.

$$(3) \quad ds^2 = c^2 \left(1 - \frac{\alpha}{R} \right) dt^2 - \frac{dR^2}{1 - \frac{\alpha}{R}} - R^2(d\theta^2 + \sin^2 \theta d\varphi^2),$$

where α is a constant depending on the mass at the origin. Levi-Civita* has given three solutions of the equations $G_{ik} = 0$, one of which includes the above, and Weyl† has given still another solution. Later Kottler‡ obtained the form (3) not by the solution of the equations $G_{ik} = 0$ but as a consequence of certain postulates. It is the purpose of this paper to accomplish the same result by the following set of postulates:

I. Assumptions *A* and *B* of Einstein, in accordance with which we write (1) in the form

$$(4) \quad ds^2 = V^2 dt^2 - ds_0^2,$$

where

$$(5) \quad ds_0^2 = \sum_{i,k}^{1,2,3} a_{ik} dx_i dx_k,$$

the functions V and a_{ik} being independent of t .

II. The function V is a solution of

$$\Delta_2 \theta = 0,$$

where $\Delta_2 \theta$ is the Beltrami differential parameter formed with respect to the form (5), and is defined by

$$(6) \quad \Delta_2 \theta = \frac{1}{\sqrt{a}} \sum_i \frac{\partial}{\partial x_i} \left(\sum_k a^{(ik)} \sqrt{a} \frac{\partial \theta}{\partial x_k} \right),$$

where a is the determinant of the functions a_{ik} and $a^{(ik)}$ is the cofactor of a in this determinant divided by a .§ This assumption is equivalent to the equation $G_{44} = 0$. In this equation and hereafter \sum_j means the sum for $j = 1, 2, 3$.

III. The surfaces $V = \text{const.}$ form part of a triply orthogonal system in the space, S_3 , of coördinates x_1, x_2, x_3 .

IV. The orthogonal trajectories of $V = \text{const.}$ in S_3 are paths of the particle, in the sense that the coördinates x_1, x_2, x_3 , of a world-line determine a path in S_3 of a particle in the gravitational field for which the world-line is the representation in terms of space and time t .

V. The form (5) is euclidean to a first approximation.

2. **Geodesics in the four-space and in S_3 .** It can be shown|| that in any three-space there exist triply-orthogonal systems of surfaces, and accord-

* Rendiconti dei Lincei, ser. 5, vol. 27 (1918), p. 365.

† Ann. der Physik, vol. 54 (1917), p. 117.

‡ Ann. der Physik, vol. 56 (1918), p. 401.

§ Bianchi, Lezioni di Geometria Differenziale, 2d ed., vol. 1, p. 68.

|| Wright, Invariants of Quadratic Differential Forms, Cambridge Tract No. 9, pp. 64-67.

ingly (5) can be given the form

$$(7) \quad ds_0^2 = \sum_i a_i dx_i^2,$$

where now the coördinate surfaces form a triply orthogonal system.

If we take s for the parameter along a world-line, and put $\dot{x}_i = dx_i/ds$, $\dot{t} = dt/ds$, the integral (2) becomes

$$(8) \quad \int \sqrt{V^2 \dot{t}^2 - \sum a_i \dot{x}_i^2} ds \equiv \int \varphi(V, x_1, x_2, x_3, \dot{x}_1, \dot{x}_2, \dot{x}_3, \dot{t}) ds.$$

The Euler equations of condition that (8) be stationary are

$$(9) \quad \frac{\partial \varphi}{\partial x_i} - \frac{d}{ds} \left(\frac{\partial \varphi}{\partial \dot{x}_i} \right) = 0^* \quad (i = 1, 2, 3, 4; x_4 = t).$$

Applying these conditions to (8), and noting that in consequence of the choice of s as parameter, we have $\varphi = 1$ along a world-line, we obtain

$$(10) \quad \frac{d^2 x_i}{ds^2} + \sum_j \frac{\partial \log a_i}{\partial x_j} \frac{dx_i}{ds} \frac{dx_j}{ds} - \frac{1}{2a_i} \sum_j \frac{\partial a_j}{\partial x_i} \left(\frac{dx_j}{ds} \right)^2 + \frac{V}{a_i} \frac{\partial V}{\partial x_i} \left(\frac{dt}{ds} \right)^2 = 0,$$

$$(11) \quad \frac{dt}{ds} = \frac{k^2}{V^2},$$

where k is a constant.

By definition the geodesics in S_3 are the curves along which the integral $\int \sqrt{\sum a_i dx_i^2}$ is stationary. When s_0 is taken for the parameter along such a geodesic, we find that the equations of a geodesic are

$$(12) \quad \frac{d^2 x_i}{ds_0^2} + \sum_j \frac{\partial \log a_i}{\partial x_j} \frac{dx_i}{ds_0} \frac{dx_j}{ds_0} - \frac{1}{2a_i} \sum_j \frac{\partial a_j}{\partial x_i} \left(\frac{dx_j}{ds_0} \right)^2 = 0.$$

From (4) and (11) it follows that the parameters s and s_0 along a world-line and the corresponding path in S_3 are in the relation

$$(13) \quad ds_0 = \sqrt{\frac{k^2}{V^2} - 1} ds.$$

When we express equations (10) in terms of s_0 , we obtain

$$(14) \quad \begin{aligned} \frac{d^2 x_i}{ds_0^2} + \sum_j \frac{\partial \log a_i}{\partial x_j} \frac{dx_i}{ds_0} \frac{dx_j}{ds_0} - \frac{1}{2a_i} \sum_j \frac{\partial a_j}{\partial x_i} \left(\frac{dx_j}{ds_0} \right)^2 \\ = \frac{k^2}{V(k^2 - V^2)} \left(\frac{dV}{ds_0} \frac{dx_i}{ds_0} - \frac{1}{a_i} \frac{\partial V}{\partial x_i} \right). \end{aligned}$$

From this equation and (12) it follows that a necessary condition that the path of a particle be a geodesic in S_3 is

* Bolza, Lectures on the Calculus of Variations, p. 123.

$$(15) \quad \frac{dV}{ds_0} \frac{dx_i}{ds_0} = \frac{1}{a_i} \frac{\partial V}{\partial x_i} \quad (i = 1, 2, 3).$$

If we multiply these respective equations by $\sqrt{a_i}$, square the resulting equations and add them, we get

$$(16) \quad \left(\frac{dV}{ds_0} \right)^2 = \sum_i \frac{1}{a_i} \left(\frac{\partial V}{\partial x_i} \right)^2 \equiv \Delta_1 V,$$

where $\Delta_1 \theta$ is the first differential parameter of θ with respect to the form (7). When ds_0^2 is written in the general form (5), the expression for $\Delta_1 \theta$ is

$$(17) \quad \Delta_1 \theta = \sum_{ik} a^{(ik)} \frac{\partial V}{\partial x_i} \frac{\partial V}{\partial x_k}.*$$

Hence (15) may be written in the form

$$(18) \quad \frac{dx_i}{ds_0} = \frac{1}{\sqrt{\Delta_1 V}} \cdot \frac{1}{a_i} \frac{\partial V}{\partial x_i}.$$

The direction of the tangent to any curve on a surface $V = \text{const.}$ through the point P where a path curve meets the surface is given by the values of dx_i/ds_1 , s_1 being the arc along the curve, where

$$(19) \quad \sum_i \frac{\partial V}{\partial x_i} \frac{dx_i}{ds_1} = 0.$$

From (18) and (19) it follows that

$$(20) \quad \sum a_i \frac{dx_i}{ds_0} \frac{dx_i}{ds_1} = 0,$$

which is the condition that the path is orthogonal to the surface, since this condition is satisfied by every curve through P .†

Conversely, any orthogonal trajectory of the surfaces $V = \text{const.}$ is defined by (18). In fact from (20) and the equation

$$\sum a_i \frac{dx_i}{ds_0} \frac{dx_i}{ds_2} = 0,$$

for a second curve on $V = \text{const.}$ we get

$$(21) \quad \frac{\frac{dx_1}{ds_0}}{a_2 a_3 \left(\frac{dx_2}{ds_1} \frac{dx_3}{ds_2} - \frac{dx_3}{ds_1} \frac{dx_2}{ds_2} \right)} = \dots = \frac{\frac{dx_3}{ds_0}}{a_1 a_2 \left(\frac{dx_1}{ds_1} \frac{dx_2}{ds_2} - \frac{dx_2}{ds_1} \frac{dx_1}{ds_2} \right)} = R,$$

* Bianchi, l. c., p. 61.

† Bianchi, l. c., p. 330.

where by composition we find

$$R = \frac{1}{\sqrt{a_1 a_2 a_3} Q}, \quad Q = \sqrt{\Sigma a_2 a_3 \left(\frac{dx_2}{ds_1} \frac{dx_3}{ds_2} - \frac{dx_3}{ds_1} \frac{dx_2}{ds_2} \right)^2}.$$

In like manner from (19) and

$$\Sigma \frac{\partial V}{\partial x_i} \frac{dx_i}{ds_2} = 0,$$

we obtain

$$(22) \quad \frac{\frac{\partial V}{\partial x_1}}{\frac{dx_2}{ds_1} \frac{dx_3}{ds_2} - \frac{dx_3}{ds_1} \frac{dx_2}{ds_2}} = \dots = \frac{\frac{\partial V}{\partial x_3}}{\frac{dx_1}{ds_1} \frac{dx_2}{ds_2} - \frac{dx_2}{ds_1} \frac{dx_1}{ds_2}} = \frac{\sqrt{a_1 a_2 a_3} \sqrt{\Delta_1 V}}{Q}.$$

From (21) and (22) follows (18).

The above results may be stated as follows:

If the path of a particle in a permanent gravitational field is a geodesic, it is an orthogonal trajectory of the surfaces $V = \text{const.}$

3. Condition that the orthogonal trajectories of the surfaces $V = \text{const.}$ be geodesics. In establishing this condition we make use of the mixed differential parameter of the first order, $\Delta_1(\theta, \varphi)$, which when formed with respect to (5) is defined by

$$\Delta_1(\theta, \varphi) = \sum_{i,k} a^{(ik)} \frac{\partial \theta}{\partial x_i} \frac{\partial \varphi}{\partial x_k}.$$

For the form (7) this is

$$\Delta_1(\theta, \varphi) = \sum_i \frac{1}{a_i} \frac{\partial \theta}{\partial x_i} \frac{\partial \varphi}{\partial x_i}.$$

From (18) we have

$$\frac{d^2 x_i}{ds_0^2} = \sum_j \frac{\partial}{\partial x_j} \left(\frac{1}{\sqrt{\Delta_1 V}} \frac{1}{a_i} \frac{\partial V}{\partial x_i} \right) \cdot \frac{1}{\sqrt{\Delta_1 V}} \frac{1}{a_j} \frac{\partial V}{\partial x_j} = \frac{1}{\sqrt{\Delta_1 V}} \Delta_1 \left(V, \frac{1}{\sqrt{\Delta_1 V}} \frac{1}{a_i} \frac{\partial V}{\partial x_i} \right).$$

Substituting in (12), we get

$$a_i \sqrt{\Delta_1 V} \Delta_1 \left(V, \frac{1}{\sqrt{\Delta_1 V}} \frac{1}{a_i} \frac{\partial V}{\partial x_i} \right) \sum_j \frac{1}{a_j} \frac{\partial \log a_i}{\partial x_j} \frac{\partial V}{\partial x_i} \frac{\partial V}{\partial x_j} - \frac{1}{2} \sum_j \frac{1}{a_j} \frac{\partial \log a_j}{\partial x_i} \left(\frac{\partial V}{\partial x_j} \right)^2 = 0,$$

which may be written

$$(23) \quad \Delta_1 \left(V, \frac{\partial V}{\partial x_i} \right) - \frac{1}{2} \frac{\partial V}{\partial x_i} \frac{\Delta_1(V, \Delta_1 V)}{\Delta_1 V} - \frac{1}{2} \sum_j \frac{1}{a_j} \frac{\partial \log a_j}{\partial x_i} \left(\frac{\partial V}{\partial x_j} \right)^2 = 0,$$

since

* Bianchi, Lezioni, p. 61.

$$(24) \quad \Delta_1(V, \Delta_1 V) = \sum_j \frac{2}{a_j} \Delta_1 \left(V, \frac{\partial V}{\partial x_j} \right) \frac{\partial V}{\partial x_j} + \sum_j \left(\frac{\partial V}{\partial x_j} \right)^2 \Delta_1 \left(V, \frac{1}{a_j} \right).$$

Since

$$\frac{1}{2} \frac{\partial}{\partial x_i} \Delta_1 V = \Delta_1 \left(V, \frac{\partial V}{\partial x_i} \right) - \frac{1}{2} \sum_j \frac{1}{a_j} \frac{\partial \log a_j}{\partial x_i} \left(\frac{\partial V}{\partial x_j} \right)^2,$$

equation (23) may be written

$$\frac{1}{2} \frac{\partial}{\partial x_i} (\Delta_1 V)^2 = \Delta_1(V, \Delta_1 V) \frac{\partial V}{\partial x_i}.$$

Consequently $\Delta_1(V, \Delta_1 V)$, and also $\Delta_1 V$, must be functions of V . But when $\Delta_1 V$ is a function of V so also is $\Delta_1(V, \Delta_1 V)$. Hence we have the theorem:

A necessary and sufficient condition that the orthogonal trajectories of the surfaces $V = \text{const.}$ be geodesics is that ΔV be a function of V .

In this case the surfaces $V = \text{const.}$ are said to form a *geodesically parallel family*.*

4. The path of a ray of light. The function V is interpreted as the velocity of light in the field, and consequently along a world-line of a ray of light $ds = 0$, as follows from (4). In order to obtain the equations of these world-lines, we apply the Fermat principle that $\int dt$ be stationary along such a line, that is the integral $\int \sqrt{V^{-2} \Sigma a_i dx_i^2}$. This gives the equations

$$\frac{d^2 x_i}{dt^2} + \frac{dx_i}{dt} \sum_j \frac{\partial}{\partial x_j} \log \frac{a_i}{V^2} \frac{dx_j}{dt} - \frac{V^2}{2a_i} \sum_j \frac{\partial}{\partial x_i} \frac{a_j}{V^2} \left(\frac{dx_j}{dt} \right)^2 = 0.$$

When we require that a path of light be a geodesic in S_3 , we obtain (18). Hence:

When the orthogonal trajectories of the surfaces $V = \text{const.}$ are paths of a particle in a permanent gravitational field, they are also the paths of a ray of light, and conversely.

5. Certain triply orthogonal systems in euclidean space. A necessary and sufficient condition that

$$ds^2 = H_1^2 dx_1^2 + H_2^2 dx_2^2 + H_3^2 dx_3^2$$

the linear element of euclidean space is that the functions H_i satisfy the following equations of Lamé:†

* For other proofs of this theorem the reader is referred to Bianchi, l. c., p. 338; also, Wright, l. c., p. 64.

† Eisenhart, Differential Geometry, p. 449.

$$(25) \quad \frac{\partial^2 H_i}{\partial x_j \partial x_k} = \frac{1}{H_j} \frac{\partial H_j}{\partial x_k} \frac{\partial H_i}{\partial x_j} + \frac{1}{H_k} \frac{\partial H_k}{\partial x_j} \frac{\partial H_i}{\partial x_k},$$

$$\frac{\partial}{\partial x_i} \left(\frac{1}{H_i} \frac{\partial H_j}{\partial x_i} \right) + \frac{\partial}{\partial x_j} \left(\frac{1}{H_j} \frac{\partial H_i}{\partial x_j} \right) + \frac{1}{H_k^2} \frac{\partial H_i}{\partial x_k} \frac{\partial H_j}{\partial x_k} = 0,$$

where $i \neq j$, $i \neq k$, $j \neq k$.

We consider the case where H_1 is a function of x_1 alone, and write

$$(26) \quad H_1 = X_1',$$

the accent indicating differentiation with respect to x_1 . Now equations (25) reduce to

$$(27) \quad \frac{\partial}{\partial x_1} \left(\frac{1}{H_3} \frac{\partial H_2}{\partial x_3} \right) = 0, \quad \frac{\partial}{\partial x_1} \left(\frac{1}{H_2} \frac{\partial H_3}{\partial x_2} \right) = 0,$$

$$(28) \quad \frac{\partial}{\partial x_1} \left(\frac{1}{H_1} \frac{\partial H_2}{\partial x_1} \right) = 0, \quad \frac{\partial}{\partial x_1} \left(\frac{1}{H_1} \frac{\partial H_3}{\partial x_1} \right) = 0,$$

$$(29) \quad \frac{\partial}{\partial x_2} \left(\frac{1}{H_2} \frac{\partial H_3}{\partial x_2} \right) + \frac{\partial}{\partial x_3} \left(\frac{1}{H_3} \frac{\partial H_2}{\partial x_3} \right) + \frac{1}{H_1^2} \frac{\partial H_2}{\partial x_1} \frac{\partial H_3}{\partial x_1} = 0.$$

From (28) we have by integration

$$(30) \quad H_2 = \sigma X_1 + \bar{\sigma}, \quad H_3 = \tau X_1 + \bar{\tau},$$

where σ , $\bar{\sigma}$, τ and $\bar{\tau}$ are independent of x_1 . In accordance with (27) these functions must satisfy the conditions

$$\tau \frac{\partial \bar{\sigma}}{\partial x_3} - \bar{\tau} \frac{\partial \sigma}{\partial x_3} = 0, \quad \sigma \frac{\partial \bar{\tau}}{\partial x_2} - \bar{\sigma} \frac{\partial \tau}{\partial x_2} = 0.$$

If we replace these equations by

$$(31) \quad \frac{\partial \bar{\sigma}}{\partial x_3} = \lambda \bar{\tau}, \quad \frac{\partial \sigma}{\partial x_3} = \lambda \tau,$$

$$\frac{\partial \bar{\tau}}{\partial x_2} = \mu \bar{\sigma}, \quad \frac{\partial \tau}{\partial x_2} = \mu \sigma,$$

we have

$$\frac{1}{H_3} \frac{\partial H_2}{\partial x_3} = \lambda = \frac{1}{\tau} \frac{\partial \sigma}{\partial x_3}, \quad \frac{1}{H_2} \frac{\partial H_3}{\partial x_2} = \mu = \frac{1}{\sigma} \frac{\partial \tau}{\partial x_2},$$

so that (29) becomes

$$(32) \quad \frac{\partial}{\partial x_2} \left(\frac{1}{\sigma} \frac{\partial \tau}{\partial x_2} \right) + \frac{\partial}{\partial x_3} \left(\frac{1}{\tau} \frac{\partial \sigma}{\partial x_3} \right) + \sigma \tau = 0.$$

Each solution of this equation determines λ and μ , and then $\bar{\sigma}$ and $\bar{\tau}$ follow from (31).

Equation (32) expresses the condition that

$$\sigma^2 dx_2^2 + \tau^2 dx_3^2$$

is the linear element of the unit sphere.*

If it is required further that $H_2 H_3$ be the product of a function of x_1 alone, say φ , and a function ψ independent of x_1 , we have, from (30)

$$\frac{\sigma\tau}{\psi} X_1^2 + \frac{\sigma\bar{\tau} + \bar{\sigma}\tau}{\psi} X_1 + \frac{\bar{\sigma}\bar{\tau}}{\psi} = \varphi.$$

From the two equations obtained by differentiating this equation with respect to x_2 and x_3 and (31) we find that $\sigma/\bar{\sigma} = \tau/\bar{\tau} = k$, where k is a constants. Consequently the general solution may be written

$$(33) \quad H_2 = X_1 \sigma, \quad H_3 = X_1 \tau.$$

6. Derivation of the Schwarzschild form (3). In accordance with postulate III, we take the surfaces $V = \text{const.}$ for the coördinate surfaces $x_1 = \text{const.}$ of a triply orthogonal system in S_3 , and write

$$(34) \quad V^2 = c^2(1 + 2\varphi_1(x_1)),$$

where c is the constant velocity of light.

From the results of §§ 2, 3 it follows that postulate IV is equivalent to $\Delta_1 V = \varphi(x_1)$, which reduces to

$$(35) \quad a_1 = \frac{V'^2}{\varphi} = \frac{c^2 \varphi_1'^2}{\varphi(1 + 2\varphi_1)}.$$

Since by postulate II we have $\Delta_2 V = 0$, we must have also

$$(36) \quad a_2 a_3 = \frac{\psi^2}{\varphi},$$

where ψ is independent of x_1 .

From the results of § 5 it follows that the linear element of euclidean space, satisfying the conditions that a_1 is a function of x_1 alone and (36), can be given the form

$$(37) \quad d\bar{s}^2 = dx_1^2 + x_1^2(\sigma^2 dx_2^2 + \tau^2 dx_3^2).$$

In accordance with postulate V the linear element of S_3 is to be (37) to a first approximation. From (35) and (36) we find that such an approximation is given by taking

$$(38) \quad \varphi_1'^2 = \frac{\varphi}{c^2}, \quad \varphi = \frac{m^2}{4x_1^4},$$

* Eisenhart, Differential Geometry, p. 157.

where m is a constant, that is $\varphi_1 = -m/2cx_1$. Then from (34), (35) and (37) we have

$$(39) \quad ds^2 = c^2 \left(1 - \frac{m}{cx_1} \right) dt^2 - \frac{1}{1 - \frac{m}{cx_1}} dx_1^2 - x_1^2 (\sigma^2 dx_2^2 + \tau^2 dx_3^2).$$

When we take the solutions $\sigma = 1$, $\tau = \sin x_2$ of (32), we obtain (3).

7. Derivation of a form due to Levi-Civita. If in (26) and (33) we put $X_1 = 1/\sqrt{K_0\mu}x_1$, where K_0 and μ are constants, we have in place of (37)

$$d\bar{s}^2 = \frac{1}{K_0\mu x_1^4} dx_1^2 + \frac{1}{x_1^2} \left(\frac{\sigma^2 dx_2^2 + \tau^2 dx_3^2}{K_0\mu} \right),$$

and the expression in parenthesis has curvature $K_0\mu$. In place of (34) we put

$$V^2 = \mu V_0^2 (1 + 2\varphi_1(x_1)),$$

where V_0 is a constant. Proceeding as in the preceding section we have in place of (38)

$$\frac{V_0^2 \varphi_1'^2}{\varphi} = \frac{1}{K_0 x_1^4 \mu^2}, \quad \varphi = m x_1^4,$$

where m is a constant. If we take accordingly

$$2\varphi_1 = -\frac{\epsilon x_1}{\mu}, \quad m = \frac{V_0^2 K_0}{4}, \quad \epsilon = \pm 1,$$

we obtain the form

$$ds^2 = V_0^2 (\mu - \epsilon x_1) dt^2 - \frac{dx_1^2}{K_0 x_1^4 (\mu - \epsilon x_1)} - \frac{1}{x_1^2} \left(\frac{\sigma^2 dx_2^2 + \tau^2 dx_3^2}{K_0 \mu} \right),$$

due to Levi-Civita.*

* L. c.